

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

Original Article

Expanded classification for immediate molar implant placement: Integrating root socket strategies beyond septal bone typing

Chiun-Lin Steven Liu ^{a,b,c,d}, Shih-Cheng Wen ^e,
Ching-Jung Chang Chien ^f, Je-Kang Du ^{b,g*}

^a School of Dental Medicine, University of Pennsylvania, Philadelphia, PA, USA

^b School of Dentistry, College of Dental Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

^c Main Line Dental Implant Center, Private Practice, Berwyn, PA, USA

^d Dr. Morton Amsterdam Museum, PA, USA

^e School of Dentistry, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan

^f Kande Dental Clinic Private Practice, Taipei, Taiwan

^g Department of Dentistry, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

Received 18 December 2025; Final revision received 6 January 2026

Available online ■ ■ ■

KEYWORDS

Classification system;
Immediate implant
placement;
Molar extraction;
Socket preservation;
Septal bone;
Treatment algorithm

Abstract *Background/purpose:* This conceptual development study with exploratory clinical analysis aimed to create and preliminarily evaluate a six-category classification system for immediate implant placement (IIP) in molar extraction sites. IIP in molar regions remains challenging due to complex socket morphology and variable septal bone availability. Existing septal-based classifications (e.g., Smith-Tarnow) may exclude viable cases with favorable root socket anatomy. The system was developed through literature review, CBCT-based anatomical analysis, and Delphi expert consensus, with clinical applicability assessed via retrospective review of 64 M IIP cases.

Materials and methods: A classification system combining septal bone conditions (Types A-C) and root socket-based anchorage (Types D, M, and P) was developed through literature review, CBCT-based anatomical assessment, retrospective case evaluation, and expert consensus via a Delphi process.

Results: Type A demonstrated highest predictability (insertion torque >30 N/cm, survival >90%). Type B commonly required grafting for gaps >2 mm, while Type C relied on adequate buccolingual cortical engagement. Types D and M provided reliable anchorage in suitable sockets with careful prosthetic planning. Type P showed consistently high primary stability (ISQ >60), minimal early marginal bone loss (<0.2 mm at 6–24 months), and favorable short-term outcomes, supporting its role as a sinus augmentation alternative in selected maxillary molars.

* Corresponding author. School of Dentistry, Kaohsiung Medical University, No.100, Shih-Chuan 1st Road, Kaohsiung 80708, Taiwan.
E-mail address: dujekang@gmail.com (J.-K. Du).

<https://doi.org/10.1016/j.jds.2026.01.011>

1991-7902/© 2026 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Please cite this article as: C.-L. Steven Liu, S.-C. Wen, C.-J. Chang Chien et al., Expanded classification for immediate molar implant placement: Integrating root socket strategies beyond septal bone typing, Journal of Dental Sciences, <https://doi.org/10.1016/j.jds.2026.01.011>

Conclusion: The proposed six-category system provides a comprehensive anatomy-based framework for molar IIP, integrating septal and root-specific considerations to expand case selection, shorten treatment time, and maintain predictable outcomes, serving as a decision-making adjunct to comprehensive anatomical and prosthetic assessment.

© 2026 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Immediate implant placement (IIP) in molar extraction sites has shown survival rates broadly comparable to delayed protocols, although its predictability remains influenced by anatomical constraints, socket morphology, and the ability to achieve adequate primary stability.^{1–5} The Smith-Tarnow classification, which categorizes molar sockets according to septal bone conditions (Types A-C), has long served as a practical reference; however, it may be less applicable in cases where septal bone is reduced or absent.^{6–8}

Recent clinical reports suggest that individual root sockets—particularly the palatal root in maxillary molars—may provide favorable anchorage and adequate stability for immediate placement, even when septal support is limited.^{9–12} These findings indicate that a broader assessment incorporating root-specific anatomy may help clinicians identify additional cases that could be managed with IIP, although further evidence is still needed.¹³

Existing classification frameworks, including those based on healing timing or bone quality, contribute valuable guidance but do not specifically address the anatomical variability within multirrooted molar sites.^{1,14,15} Treating the molar socket as a single unit may therefore overlook distinct morphological features that could influence implant selection and placement strategy.¹⁶

Although septal-based systems such as the Smith–Tarnow classification have guided clinical decision-making for over a decade, they remain fundamentally septum-dependent and present several clinically significant limitations. These systems do not fully account for root-specific anatomy (mesial, distal, or palatal sockets), apical bone availability, or prosthetically driven implant trajectory in multirrooted molar sites. Consequently, molars with compromised or absent septa are frequently directed toward delayed protocols or require ultra-wide fixtures, even when individual root sockets could provide adequate primary stability with more favorable biomechanics and safer distances from vital structures such as the maxillary sinus or inferior alveolar nerve.¹⁷

Additionally, current septal-only approaches offer limited guidance for borderline or mixed-morphology molars, where residual septal bone coexists with potentially useable individual root sockets, creating diagnostic ambiguity and inconsistent case selection. These gaps highlight the need for a more comprehensive framework that integrates both septal bone conditions and root socket-specific strategies.

In response to these limitations, the present study introduces a six-category classification that combines traditional septal bone-based types (A-C) with three additional categories reflecting individual root socket utilization (Types D, M, and P). This expanded system is intended to provide a more comprehensive, anatomy-oriented framework for IIP planning in molar regions, with the aim of supporting more consistent and predictable clinical decision-making.

Accordingly, this study was designed as a combined conceptual development and exploratory clinical applicability investigation, rather than a formal outcome-based validation trial. The study aimed to develop a six-category, anatomy-based classification system for immediate implant placement (IIP) in molar extraction sites through literature review, CBCT-based anatomical analysis, and expert consensus, and to assess its initial clinical feasibility and applicability via retrospective evaluation of 64 M IIP cases using predefined diagnostic criteria.

Materials and methods

Study design

This conceptual development study with exploratory analysis aimed to create a six-category classification system for molar IIP through systematic literature synthesis, CBCT-based anatomical analysis, and expert consensus (development phase), followed by retrospective evaluation of 64 clinical cases to test clinical applicability (exploratory assessment phase).

Development of preliminary classification categories (literature review and CBCT-based anatomical analysis)

A six-type classification for IIP in molar extraction sites was developed through an integrative process combining a structured literature review, CBCT-based anatomical analysis, retrospective clinical pattern evaluation, and expert consensus. The primary objective was to establish an anatomy-driven framework reflecting both septal bone conditions and individual root socket morphology.

Literature review

A targeted search of PubMed and MEDLINE databases was performed for peer-reviewed articles published between

2000 and 2023 using the keywords “immediate implant placement,” “molar extraction,” “septal bone,” “root socket anatomy,” and “primary stability.” Studies were screened based on relevance to molar socket morphology, implant stability, and clinical decision-making for IIP, with emphasis on longitudinal studies and systematic reviews.^{18,19}

CBCT-based anatomical analysis

Representative anonymized CBCT datasets of molar extraction sites were analyzed using specialized planning software (Blue Sky Plan, Blue Sky Bio LLC, Libertyville, IL, USA) to characterize socket morphology, including buccolingual width, presence of the interradicular septum, root divergence, apical bone height, and proximity to the maxillary sinus or inferior alveolar nerve.^{20,21} Intra-observer reliability for quantitative CBCT measurements (septal height/width, root divergence angles, apical bone height, anatomical clearance) was assessed by repeating analysis on 20 % of cases ($n = 13$), yielding an intra-class correlation coefficient (ICC) of 0.92 (95 % CI: 0.85–0.97), confirming measurement consistency.

Clinical case exploratory analysis (retrospective analysis of 64 M IIP cases)

A retrospective review was conducted on 64 consecutive cases of immediate implant placement in molar extraction sites (38 maxillary, 26 mandibular) treated at the Department of Dentistry, Kaohsiung Medical University Hospital, between January 2018 and December 2023.

Inclusion criteria:

1. Patients who received immediate implant placement in single molar extraction sites (maxillary or mandibular).
2. Availability of pre-operative and post-operative CBCT imaging suitable for assessment of socket morphology and anatomical constraints.
3. Minimum 12-month clinical and radiographic follow-up after functional loading.
4. Documentation of primary stability parameters at placement (insertion torque and/or ISQ), prosthetic restoration type, and peri-implant bone levels.

Exclusion criteria:

1. Multiple adjacent implants involving segmental edentulism or full-arch rehabilitation.
2. Sites with acute uncontrolled infection, untreated periodontal disease, or non-restorable vertical root fractures extending into the furcation.
3. Patients with uncontrolled systemic conditions (e.g., uncontrolled diabetes), history of head and neck radiotherapy, or medications significantly affecting bone metabolism (e.g., high-risk antiresorptives).
4. Heavy smokers (≥ 20 cigarettes/day) or patients with inadequate records preventing reliable assessment of outcomes.

5. Cases with follow-up shorter than 12 months or incomplete radiographic documentation.

Primary stability assessment and gap management

Insertion torque (IT) was recorded using a manual torque wrench during implant placement (N/cm). Resonance frequency analysis (RFA) was performed immediately post-placement and at 4 weeks using Osstell Beacon (Integration Diagnostics AB, Sweden), with ISQ values reported as mean \pm SD. Gap management was tailored to socket morphology: Type A/B (self-contained, no graft); Type C/D/M/P (containment defects, xenograft granules \pm resorbable membrane).

Follow-up and radiographic evaluation

Periapical radiographs were obtained immediately post-placement, at 4 weeks (prosthetic conversion), 3 months, and 12 months. Marginal bone level (MBL) changes were measured from implant platform to first bone-to-implant contact (mesial/distal) using ImageJ software, calibrated against known implant dimensions. MBL change was calculated as $\Delta\text{MBL} = \text{MBL}_{\text{follow-up}} - \text{MBL}_{\text{baseline}}$ (mm).

Protocol characteristics

This retrospective analysis reflects real-world clinical practice rather than standardized prospective protocols. Key characteristics across the 64 cases were:

- Implant systems: Astra Tech® (Osseospeed™ TX; $n = 37$, 58 %) and Osstem® (Hiossen; $n = 27$, 42 %); both moderately rough hydrophilic surfaces (TiUnite®/SLActive®).
- Dimensions: Platform diameters ranged from 5.0 to 7.0 mm, and lengths from 10 to 13 mm. Platform diameter refers to implant neck diameter at the implant–abutment interface, with a cohort mean of 5.4 ± 0.4 mm, reflecting clinical selection based on socket morphology, bone quality, and prosthetic requirements.
- Surgical approach: Flapless ($n = 59$, 92 %); root sectioning for all multi-rooted sites; primary stability threshold ≥ 30 N/cm or ISQ ≥ 60 required for IIP.
- Grafting: Xenograft (Bio-Oss®, Geistlich; $n = 44$, 68 %), allograft (cortical/cancellous 50:50; $n = 16$, 25 %), no graft (small gaps < 2 mm; $n = 4$, 7 %).
- Loading: Immediate provisionalization (non-occlusal; $n = 18$, 28 %), early loading (6–8 weeks; $n = 33$, 52 %), conventional (3–4 months; $n = 13$, 20 %).

Protocol distribution by classification type is summarized in Table 1 ($\chi^2 P = 0.42$, no significant bias).

Data extracted included implant stability at insertion (torque and ISQ), grafting procedures, prosthetic restoration type, and radiographic bone maintenance. All cases met the inclusion criteria for IIP following molar extraction and were evaluated for implant site characteristics, root socket utilization, and primary stability parameters.^{22,23} All data were de-identified prior to analysis, and the protocol

Table 1 Protocol characteristics by classification type (n = 64).

Type	Astra Tech® Osseospeed™ TX n (%)	Osstem® Hiossen n (%)	Platform diameter (mm)	Xenograft n (%)	Early/immediate loading n (%)
A (n = 18)	11 (61 %)	7 (39 %)	5.2 ± 0.3	10 (56 %)	12 (67 %)
B (n = 15)	9 (60 %)	6 (40 %)	5.4 ± 0.4	12 (80 %)	9 (60 %)
C (n = 8)	5 (63 %)	3 (37 %)	5.7 ± 0.3	7 (88 %)	4 (50 %)
D (n = 9)	5 (56 %)	4 (44 %)	5.3 ± 0.4	6 (67 %)	6 (67 %)
M (n = 7)	4 (57 %)	3 (43 %)	5.2 ± 0.3	4 (57 %)	4 (57 %)
P (n = 7)	3 (43 %)	4 (57 %)	5.1 ± 0.3	5 (71 %)	5 (71 %)
Total	37 (58 %)	27 (42 %)	5.4 ± 0.4	44 (69 %)	40 (63 %)

χ^2 test for protocol distribution: $P = 0.42$ (no significant bias).

was approved by the Institutional Review Board (IRB No. KMUHIRB-E(I)-20250331).

Expert consensus and final classification (three-round Delphi process with $\geq 80\%$ agreement across five diagnostic domains)

Preliminary classification categories were formed based on anatomical patterns and clinical outcomes identified in the literature and case review. Five evaluative domains—socket morphology, primary stability, anatomical safety, prosthetic feasibility, and long-term radiographic performance were defined. Preliminary categories were reviewed through a three-round Delphi process involving 6 specialists (3 implantologists, 2 prosthodontists, 1 oral radiologist; each with ≥ 10 years implant experience). Inter-observer reliability testing: Before Delphi rounds, Kappa analysis was conducted on 15 representative CBCT cases (initial $\kappa = 0.78$, 95 % CI: 0.62–0.94; post-training/clarification

$\kappa = 0.92$, 95 % CI: 0.81–0.98). Consensus was defined as $\geq 80\%$ agreement;²⁴ final case assignments achieved 100 % agreement across all 64 retrospective cases after round 3.

Classification criteria and definitions

The final system comprises six distinct types based on septal and root socket anchorage (Figs. 1 and 2):

- Type A: Complete septal containment.
- Type B: Partial septal support
- Type C: Peripheral socket engagement.
- Type D: Distal root socket anchorage.
- Type M: Mesial root socket anchorage.
- Type P: Palatal root socket anchorage.

Diagnostic parameters for classification

Each classification type was defined according to standardized diagnostic parameters:




Figure 1 Smith-Tarnow septal-based classification for immediate molar implant placement and treatment algorithm. Type A: complete septal containment. Type B: partial septal stabilization. Type C: peripheral socket engagement. The flowchart illustrates the decision algorithm based on interradicular septal integrity and transition to root socket–based strategies when septal support is inadequate.




Figure 2 Root socket-based immediate molar implant placement (Types D, M, and P) and study design flowchart. Type D: distal root socket placement. Type M: mesial root socket placement. Type P: palatal root socket placement, with or without sinus augmentation. The flowchart summarizes the study process, including literature review, CBCT-based anatomical analysis, retrospective case evaluation, Delphi consensus, and final classification development.

1. Socket morphology: CBCT assessment of available supporting bone; sockets with ≥ 3 mm of circumferential septal support were classified as Type A.^{25,26}
2. Primary stability: Sites were considered suitable for IIP when adequate primary stability was anticipated based on bone density, socket morphology, and preoperative CBCT assessment, typically corresponding to insertion torque values around 30 N/cm and ISQ values near or above 60, as suggested in the literature.^{4,5}
3. Anatomical safety: A minimum of 2 mm clearance from the maxillary sinus or inferior alveolar nerve was required, based on preoperative CBCT.^{6,27}
4. Prosthetic emergence profile feasibility: socket axis and angulation were assessed against prosthetic path alignment, using diagnostic wax-ups or virtual planning. Type M and D sockets required verification that implant emergence would not result in severe buccal displacement or prosthetic compromise.^{28,29}
5. Long-term predictability: Marginal bone loss < 0.5 mm over 12–24 months was used as an indicator of stability, consistent with prior literature. This benchmark supported inclusion of Types P and D in appropriate scenarios.^{18,30–33}

When multiple socket options appeared feasible, the final type assignment was based on the root socket offering the most favorable balance between anatomical safety, prosthetic alignment potential, and primary stability.

Results

Among the 64 retrospectively reviewed molar IIP cases, the distribution by classification type was as follows: Type A ($n = 12$, 19%), Type B ($n = 15$, 23%), Type C ($n = 14$,

22%), Type D ($n = 10$, 16%), Type M ($n = 8$, 12%), and Type P ($n = 5$, 8%), reflecting clinically representative prevalence within the cohort (Table 1).

The novel six-category classification system

A six-category classification system for IIP in molar extraction sites was developed to address the limitations of septal bone-only concepts. The system integrates septal bone morphology (Types A–C) with root socket-specific strategies (Types D, M and P), acknowledging that palatal, mesial, and distal sockets may provide adequate primary stability and favorable mid-term outcomes in well-selected cases. Within the retrospective cohort, Type A and Type P sites were more frequently observed in the maxilla, whereas Types B, C, D, and M were more commonly encountered in mandibular molars. This framework enables a more comprehensive anatomical assessment of molar sockets and may broaden the range of sites considered suitable for IIP. An overview of the classification types is summarized in Table 2, with representative clinical examples illustrated in Figs. 3–8.

Primary stability parameters were recorded descriptively across socket types, with a mean insertion torque of 35 ± 12 N/cm and mean ISQ values of 72 ± 8 at placement and 78 ± 6 at 4 weeks. Mean marginal bone level change at 12 months was -0.4 ± 0.3 mm.

The clinical cases and imaging figures presented (Figs. 3–8) are illustrative examples intended to demonstrate the anatomical characteristics, surgical workflow, and practical application of each classification type. These cases are provided for descriptive and educational purposes and are not intended to serve as evidentiary or outcome-based comparisons among classification categories.

Table 2 Novel six-category classification system. Comprehensive comparison of immediate implant placement types in molar extraction sites.

Novel six-category classification system							
Comprehensive comparison of immediate implant placement types in molar extraction sites							
Classification type	Definition & placement	Primary stability	Success rate	Technical difficulty	Key advantages	Main limitations	Ideal indications
Type A complete septal containment	Implant completely contained within septal bone circumferentially around platform ⁽⁸⁾	> 30 N/cm ⁵ (TV) ⁽⁵⁾ easily achieved	> 90% ⁽¹⁸⁾ highest predictability	Easy ⁽³⁷⁾ simplest technique	<ul style="list-style-type: none"> Highest predictability⁽¹⁸⁾ No grafting required^(34,35) Immediate provisionalization possible Minimal complications 	<ul style="list-style-type: none"> Limited to cases with adequate septal bone Requires careful extraction technique⁽³⁷⁾ Good bone quality 	<ul style="list-style-type: none"> Adequate septal bone height/width Intact extraction socket Good bone quality
Type B partial septal stabilization	Sufficient septal bone for stability but gaps present between implant and socket walls ⁽⁸⁾	Variable achievable ⁽⁴⁾ but inconsistent	Moderate ⁽⁸⁾ case-dependent	Moderate careful case selection	<ul style="list-style-type: none"> Gaps < 2mm fill naturally^(6,40) Flapless technique possible⁽³⁷⁾ Good outcomes with proper selection 	<ul style="list-style-type: none"> Variable primary stability May require grafting⁽³⁸⁾ Buccal wall integrity crucial⁽⁴¹⁾ Gaps assessable 	<ul style="list-style-type: none"> Some septal bone present Intact buccal wall⁽³⁷⁾ Gaps assessable
Type C peripheral socket	No septal bone available; implant engages peripheral socket walls for stability ⁽⁸⁾	Challenging wide-diameter implants required ^(7,14,45)	Variable ⁽⁸⁾ technique-sensitive	Difficult high technical demands	<ul style="list-style-type: none"> Option when septal bone absent⁽²⁰⁾ Utilizes peripheral wall thickness Suitable for specific anatomies 	<ul style="list-style-type: none"> Requires thick buccal/lingual walls⁽²⁰⁾ Wide-diameter implants needed High technical difficulty 	<ul style="list-style-type: none"> No septal bone Thick peripheral walls (> 1mm)⁽²⁰⁾ Hourglass socket morphology
Type D distal root socket	Implant placed specifically into distal root socket of extracted molar	Good potential ^(4,15) quality bone dependent	Good case-specific	Moderate ⁽¹⁴⁾ requires proper positioning	<ul style="list-style-type: none"> Utilizes strong distal root Preserves other sockets⁽⁹⁾ Suitable for both arches 	<ul style="list-style-type: none"> May require angulated abutments Emergence profile challenges⁽²⁷⁾ Distance from adjacent teeth 	<ul style="list-style-type: none"> Adequate distal root length⁽¹⁵⁾ Good bone quality⁽¹⁵⁾ Favorable angulation⁽²⁸⁾ Sufficient apical bone⁽²⁹⁾
Type M mesial root socket	Implant positioned in mesial root socket, typically the largest root in mandibular molars	Good ⁽¹⁵⁾ usually favorable	Good consistent outcomes	Moderate standard technique	<ul style="list-style-type: none"> Often largest root socket Generally favorable angulation Good bone quality typically⁽¹⁵⁾ Suitable access 	<ul style="list-style-type: none"> Proximity to adjacent teeth Angulation considerations⁽⁹⁾ Root extraction challenges 	<ul style="list-style-type: none"> Strong mesial root⁽²¹⁾ Adequate bone quality⁽¹⁵⁾ Appropriate angulation⁽²⁸⁾ Sufficient distance from adjacent teeth
Type P palatal root socket	Implant placed into palatal root socket of maxillary molars (strongest/longest root)	> 60 ISQ consistently high	100 % (0.19 mm bone loss)	Easy minimally invasive	<ul style="list-style-type: none"> Highest success rates reported Strongest root anatomy Avoids sinus complications Minimal bone loss Simplified protocol 	<ul style="list-style-type: none"> Limited to maxillary molars Requires adequate bone height Palatal access considerations 	<ul style="list-style-type: none"> ≥ 6mm bone height around palatal root Healthy adjacent tissues No acute infection Maxillary molar sites

High success (> 90 %) Moderate success Variable success

Type A: complete septal containment (Fig. 3)

Type A sockets exhibit demonstrate circumferential septal bone surrounding the implant platform,⁸ typically achieving insertion torque values 30 N/cm⁵ and survival rates exceeding 90 % in well-selected cases.¹⁸ Immediate provisionalization is feasible when adequate primary stability is obtained.^{34–36} Flapless extraction with root sectioning preserves septal integrity,³⁷ and buccal grafting is required only when facial plate thickness is limited.^{9,38}

Type B: partial septal stabilization (Fig. 4)

Type B sockets provide partial septal support and represent an intermediate morphology between Types A and C. Primary stability is generally achievable but more variable because peripheral gaps are commonly present.⁸ Small gaps (<2 mm) may heal without augmentation, whereas larger defects often require grafting and delayed loading.^{6,37,39,40} Loss of buccal plate integrity increases the risk of failure and may indicate conversion to a delayed protocol with ridge preservation.^{3,35,41}

Type C: peripheral socket engagement (Fig. 5)

Type C sockets lack sufficient septal bone and depend on engagement of peripheral buccal and lingual cortical walls for stability.⁸ Wide-diameter implants (approximately 6–7 mm) and favorable “hourglass” morphology are often required to obtain reliable cortical anchorage,²⁰ particularly in mandibular second molars where buccolingual ridge width (~9 mm) and cortical thickness are greater

(~9 mm).²⁰ These cases demand meticulous CBCT-based planning, careful implant macrodesign selection, and atraumatic site preparation to avoid fracture of thin socket walls.

Type D: distal root socket placement (Fig. 6)

Type D refers to IIP performed within the distal root socket when socket length, angulation, and apical bone volume allow stable anchorage. This approach can be applied in both maxillary and mandibular molars. Implants are positioned along a prosthetically driven path within the distal socket, and residual sockets may be grafted to maintain ridge volume and soft-tissue contours.^{9,14,42} Distal positioning may necessitate angulated abutments to achieve an acceptable emergence profile.^{29,43}

Type M: mesial root socket placement (Fig. 7)

Type M refers to implant placement in the mesial root socket, mainly in mandibular molars. The mesial socket frequently offers favorable width, depth, and central bone morphology, making it a viable option when septal support is limited and root divergence is modest.^{15,21,44–47} Adequate apical bone, sufficient distance from adjacent teeth, and a prosthetically acceptable trajectory are essential to obtain predictable primary stability.²⁹

Type P: palatal root socket placement (Fig. 8)

Type P describes implant placement in the palatal root socket of maxillary molars, utilizing the palatal root's




Figure 3 Representative clinical workflow for type A IIP in a molar extraction site. The three roots of tooth #26 were extracted individually to preserve septal integrity. A 5 × 11 mm implant was placed directly into the septal bone, achieving high primary stability. A 1:1 mixture of cortical and cancellous allograft was applied to support healing, and second-stage surgery was completed during the same session. Sequential images illustrate caries assessment, tooth sectioning, atraumatic extraction, septal evaluation, immediate implant placement, and healing abutment insertion with soft-tissue management. The final images show definitive prosthetic restoration and a 2-year radiographic follow-up confirming stable osseointegration and appropriate implant positioning. This representative case is presented for illustrative purposes to demonstrate application of the proposed classification.




Figure 4 Representative clinical and radiographic workflow for type B IIP in a molar extraction site. The molar roots of tooth #46 were extracted individually to preserve residual septal bone. An implant was placed within the partially supportive septum, followed by selective grafting and healing abutment placement during the same session. The upper images illustrate atraumatic extraction, socket debridement, implant positioning, and soft-tissue management. The lower images present sequential periapical radiographs from preoperative assessment through immediate placement to definitive restoration, demonstrating stable integration at 2-year follow-up. This representative case is presented for illustrative purposes to demonstrate application of the proposed classification.



Figure 5 Representative clinical and radiographic workflow for type C IIP in a molar extraction site without septal bone. Tooth #37 and an associated radicular cyst were removed, resulting in a socket lacking septal support. The cystic defect was grafted, followed by implant placement relying on peripheral socket engagement, with additional grafting to fill residual gaps and concurrent healing abutment placement. The upper images illustrate extraction, defect management, implant installation, and soft-tissue closure, whereas the lower images show sequential radiographs from preoperative evaluation through immediate placement to 3-year follow-up, demonstrating stable implant integration. This case is included for illustrative purposes and does not constitute outcome-based evidence.

length, dense bone, and biomechanical advantages. Contemporary reports show high insertion torque, ISQ values typically ≥ 60 , minimal marginal bone loss at 6–24 months (0.11–0.19 mm),²¹ and excellent short-term survival, while often avoiding maxillary sinus augmentation.^{10,48,49} Clinical protocols usually include flapless extraction with root sectioning,^{22,37} palatal socket preparation at a modest palatal inclination (average 9.6°),³⁷ and buccal socket grafting with Xenograft and/or allograft to support ridge contour and soft-tissue thickness.³⁸ Incorporation of Type P recognizes a minimally invasive, biologically favorable option for selected maxillary molars.

Discussion

IIP in molar sites remains challenging due to variable socket morphology, inconsistent septal bone availability, and the proximity of vital anatomical structures. Although septal-based systems such as the Smith-Tarnow classification have provided a useful reference for more than a decade,⁸ they remain fundamentally septum-dependent and do not fully incorporate root-specific anatomy, apical bone availability, or prosthetically driven implant trajectory.^{3,16} Compared with more recent CBCT- or osseodensification-based molar socket classifications, the present system emphasizes prosthetically driven socket selection rather than bone expansion or implant macrodesign alone. Consequently, many sites with reduced or absent septa may be directed

toward delayed protocols even when individual mesial, distal, or palatal root sockets could provide adequate primary stability with favorable biomechanics.⁵⁰

The six-category framework proposed in this study integrates septal bone-based types (A-C) with root socket-based strategies (D, M, P), providing a more anatomy-driven and clinically versatile approach. By incorporating three-dimensional assessment of septal morphology, root divergence, apical bone height, and anatomical limitations,⁵⁰ the system facilitates selection of the most favorable socket for IIP while maintaining a prosthetically appropriate implant trajectory. This framework may be particularly advantageous in borderline or mixed-morphology molars where residual septal bone coexists with useable individual root sockets. The corresponding treatment hierarchy-prioritizing Type A, followed by Type P in the maxilla and Types D and M when root socket anatomy is favorable, with Types B and C or delayed placement reserved for higher-risk scenarios-supports reproducible case selection, aligns well with digital planning workflows, and promotes consistency across operators. It should be emphasized that the proposed classification system is intended to function as a decision-support framework rather than a replacement for clinician judgment. Final treatment decisions should remain guided by comprehensive CBCT-based anatomical assessment, intraoperative findings, prosthetic planning, and patient-specific risk factors. Within this context, the classification is designed to assist clinicians in organizing anatomical information and




Figure 6 Representative clinical and radiographic workflow for type D IIP using the distal root socket. Two roots of tooth #36 were extracted individually, and an implant was placed within the distal root socket to achieve primary stability, followed by selective grafting and concurrent healing abutment placement. The upper images illustrate extraction, socket preparation, implant positioning, and peri-implant soft-tissue healing. The lower images present sequential radiographs from preoperative assessment through implant placement to definitive restoration at 5-year follow-up, demonstrating stable integration and functional maintenance. This representative case is presented for illustrative purposes to demonstrate application of the proposed classification.

standardizing decision-making, while individualized clinical expertise remains essential.

In selected maxillary molars, Type P placement may be considered instead of sinus augmentation when sufficient palatal root length and bone density allow predictable anchorage without violating the sinus floor. This approach is particularly beneficial when buccal residual bone height is inadequate yet palatal height remains favorable, thereby avoiding sinus elevation that would otherwise be required solely to achieve primary stability. Type P may also be advantageous in patients with systemic comorbidities, smoking habits, or sinus anatomy that increases the morbidity or complexity of sinus lift procedures. Under such conditions, palatal socket anchorage offers a minimally invasive, graft-sparing option that maintains stability while reducing procedural risk.

Nevertheless, several borderline conditions warrant staged treatment. These include sockets with insufficient apical bone height (<3–4 mm), acute infection, compromised socket walls, or marked mesial–distal root divergence preventing a prosthetically acceptable trajectory. Cases with thin but dense cortical plates at risk of fracture, or palatal sockets with marginal defects that undermine Type P anchorage, may similarly benefit from ridge preservation followed by delayed implant placement. When CBCT evaluation or intraoperative findings suggest doubtful primary stability or conflicting anatomical versus prosthetic requirements, a delayed approach remains the more predictable option.

Incorrect application of the classification may lead to clinical complications. Misuse of Type P placement in cases with insufficient palatal bone height or excessive palatal inclination may increase the risk of sinus perforation or inadequate primary stability. Inappropriate Type D positioning can compromise prosthetic emergence profile and functional loading, whereas misapplication of Type C in ridges with inadequate buccolingual width or fragile cortical plates may elevate the risk of early implant failure. These considerations underscore the importance of strict adherence to predefined anatomical and prosthetic criteria.

In the present retrospective cohort, no type-specific complication patterns were identified among the 64 included cases during the available follow-up period. No major complications—including early implant failures, sinus perforations, or excessive marginal bone loss (>2 mm at 12 months) were observed across any classification type. Type-specific contraindications and anatomical risk thresholds were predefined and strictly applied during case selection, including avoidance of Type P placement when palatal bone height was <4 mm or sinus floor contact was <1 mm, exclusion of Type D placement when the prosthetic emergence angle exceeded 25° or a buccal wall defect >4 mm was present, and contraindication of Type C placement in sites with buccolingual ridge width <6 mm (Table 3).

Although the absence of complications should not be interpreted as evidence of predictive validity, it supports the internal consistency and clinical plausibility of the




Figure 7 Representative clinical and radiographic workflow for type M IIP using the mesial root socket. The roots of tooth #36 were extracted individually, and an implant was placed within the mesial root socket selected for favorable anatomy and stability, followed by selective grafting and concurrent healing abutment placement. The upper images illustrate atraumatic extraction, socket preparation, focused mesial placement, and soft-tissue healing. The lower images present sequential radiographs from preoperative evaluation through implant placement to definitive restoration at 3-year follow-up, demonstrating stable integration. This case is included for illustrative purposes and does not constitute outcome-based evidence.

proposed contraindication framework when applied within defined anatomical limits. Systematic complication analysis and long-term safety assessment will require future prospective studies and were beyond the scope of this exploratory investigation.

A major contribution of this classification is the integration of Type P placement, which demonstrates high primary stability ($ISQ \geq 60$), minimal early marginal bone loss, and favorable short-term outcomes,^{4,10,16,48,49} consistent with its robust bone support and advantageous biomechanics. Avoidance of sinus augmentation further aligns with minimally invasive workflows and may enhance patient comfort and treatment efficiency.^{28,29}

A key strength of the system is its reliance on objective and reproducible diagnostic criteria- CBCT-based socket morphology, insertion torque thresholds, and ISQ benchmarks,^{4,5,10,14,15} and type-specific indications and contraindications-combined with a clear treatment hierarchy. These features may improve communication between surgical and restorative clinicians while reducing inappropriate case selection,^{3,11} a known contributor to early IIP failure. These features reduce ambiguity and support more consistent case selection across clinicians with different experience levels. The system's consistency with minimally invasive workflows,^{10,14-16} emphasis on socket preservation, and applicability to immediate provisionalization in appropriately selected cases further enhance its clinical practicality (Table 2). The representative clinical cases and imaging figures are included to facilitate understanding of

the classification framework and its clinical application, rather than to provide evidentiary support for outcome comparison or effectiveness assessment.

Several limitations should be acknowledged. First, much of the supporting evidence is derived from retrospective or single-center cohorts,^{44,49} and available outcome data remain largely short-to mid-term^{3-5,10,14,15,51-54} limiting generalizability and direct inter-type comparisons, particularly in the presence of heterogeneity in surgical techniques and implant systems. These factors underscore the need for multicenter prospective studies.¹³ Second, accurate application of the proposed classification requires careful CBCT interpretation and meticulous intraoperative socket assessment, rendering the system partly operator-dependent; misclassification may occur in borderline or complex mixed-morphology cases. In addition, key biomechanical and patient-related risk modifiers-such as implant macrodesign, loading strategies, bruxism, periodontal history, and occlusal patterns-are not yet fully incorporated into the current framework.

Importantly, this study should be interpreted as a conceptual development investigation with exploratory clinical applicability, rather than a formal outcome-based, predictive, or comparative validation trial. Although retrospective evaluation of 64 cases demonstrated clinical applicability across all six classification types, the study was not designed or powered to compare long-term outcomes among categories, establish survival superiority, assess predictive accuracy, or determine inter-observer



Figure 8 Representative clinical and radiographic workflow for type P IIP using the palatal root socket with CBCT guidance. The roots of tooth #26 were extracted individually, and an implant was placed within the palatal root socket selected for its favorable bone support, followed by selective grafting and concurrent healing abutment placement. The upper images illustrate socket evaluation, grafting, palatal socket-focused implant positioning, and soft-tissue healing. CBCT images confirm three-dimensional implant trajectory and surrounding bone support. The lower images present sequential radiographs from preoperative assessment through implant placement to definitive restoration at 2-year follow-up, demonstrating stable integration. This representative case is presented for illustrative purposes to demonstrate application of the proposed classification.

Table 3 Type-specific indications, contraindications, risk thresholds, and complications (n = 64 cases).

Socket type	n (%)	Primary indication	Absolute contraindications	Risk thresholds	Complications observed
A (septum-intact)	18 (28 %)	Septum height ≥ 5 mm, width ≥ 4 mm	Apical bone < 3 mm; acute infection	IT < 30 N/cm; ISQ < 60	0/12 (0 %)
B (septum-reduced)	15 (23 %)	Septum ≥ 3 mm with buccal support	Buccal wall defect > 50 %; vertical fracture	Buccal thickness < 1 mm; MBL > 1.5 mm	0/15 (0 %)
C (septum-absent)	8 (13 %)	Buccal shelf ≥ 6 mm width	BL width < 6 mm; buccal fenestration > 4 mm	Cortical fracture; IT < 35 N/cm	0/8 (0 %)
D (distal socket)	9 (14 %)	Distal root intact ≥ 6 mm; prosthetic angle $\leq 20^\circ$	Emergence angle $> 25^\circ$; distal wall > 3 mm	Compromised emergence	0/9 (0 %)
M (mesial socket)	7 (11 %)	Mesial root ≥ 6 mm, buccal support	Mesial inclination $> 30^\circ$; periodontal loss > 50 %	Loading instability	0/7 (0 %)
P (palatal socket)	7 (11 %)	Palatal root ≥ 6 mm height; sinus floor ≥ 2 mm	Sinus contact < 1 mm; palatal inclination $> 25^\circ$	Sinus perforation	0/7 (0 %)
Total	64 (100 %)				0/64 (0 %)

IT = insertion torque; ISQ = implant stability quotient; MBL = marginal bone loss; BL = buccolingual. No early failures, sinus perforations, or excessive bone loss (> 2 mm at 12 months) observed across cohort.

reproducibility. While an exploratory association between socket classification and recorded primary stability parameters was observed ($r = 0.67$, $P < 0.01$), this finding should be interpreted descriptively and does not constitute evidence of predictive or outcome-based validity. Definitive validation will require multicenter prospective studies with standardized protocols, formal reliability testing (e.g., κ statistics), and head-to-head comparisons against established classification systems to assess long-term implant survival, marginal bone maintenance, and prosthetic outcomes.

Future refinement should incorporate CBCT-driven quantitative scoring systems^{14,15} combined with AI-assisted image analysis^{55–57} to standardize socket interpretation and reduce operator variability. Advances in CBCT technology could incorporate AI-driven automatic identification of implant site classification from Type A through Type P, with real-time alerts for contraindications based on anatomical and stability criteria. Such AI integration would further streamline clinical workflow by automatically suggesting optimal implant sizing and prosthetic axis alignment, enhancing precision, reducing operator variability, and improving overall treatment predictability. Additionally, bone quality and soft tissue conditions warrant deeper integration into future iterations of the classification. Dense bone (Lekholm & Zarb I–II) appears favorable for all six types, whereas low-density bone may justify delayed placement.¹⁵ Accumulating evidence also underscores the importance of peri-implant mucogingival stability for long-term outcomes,³⁴ yet this factor is not explicitly represented in the current framework.

Overall, the proposed six-category classification provides a structured, anatomy-based approach that expands treatment possibilities, supports evidence-based decision-making, and may shorten treatment duration in appropriately selected cases. Prospective multicenter validation and incorporation of biomechanical and patient-specific risk factors will be essential to further refine the system and to define its long-term role in molar IIP planning.

Declaration of competing interest

The authors have no conflicts of interest relevant to this article.

Acknowledgements

This research was supported by the National Science and Technology Council, ROC through a grant (NSTC 114-2314-B-037-027-), Kaohsiung Medical University grant (KMU-TB114004-2) and the Kaohsiung Medical University Hospital grant (KMUH113-3M43). The funding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results. This article is dedicated to Dr. Arnold Weisgold, the late Dr. Morton Amsterdam, Dr. Walter Cohen and Dr. Jay Seibert, my mentors in Periodontics and Periodontal Prosthesis.

References

- Chen ST, Wilson Jr TG, Hämmerle CH. Immediate or early placement of implants following tooth extraction: review of biologic basis, clinical procedures, and outcomes. *Int J Oral Maxillofac Implants* 2004;19(Suppl):12–25.
- Schwartz-Arad D, Chaushu G. The ways and wherefores of immediate placement of implants into fresh extraction sites: a literature review. *J Periodontol* 1997;68:915–23.
- Esposito M, Grusovin MG, Polyzos IP, et al. Timing of implant placement after tooth extraction: immediate, immediate-delayed or delayed implants? A cochrane systematic review. *Eur J Oral Implant* 2010;3:189–205.
- Sennerby L, Meredith N. Implant stability measurements using resonance frequency analysis: biological and biomechanical aspects and clinical implications. *Periodontol* 2000 2008;47: 51–66.
- Otoni JM, Oliveira ZF, Mansini R, et al. Correlation between placement torque and survival of single-tooth implants. *Int J Oral Maxillofac Implants* 2005;20:769–76.
- Ferrus J, Cecchinato D, Pjetursson EB, et al. Factors influencing ridge alterations following immediate implant placement into extraction sockets. *Clin Oral Implants Res* 2010;21: 22–9.
- Araújo MG, Sukekava F, Wennström JL, et al. Ridge alterations following implant placement in fresh extraction sockets: an experimental study in the dog. *J Clin Periodontol* 2005;32: 645–52.
- Smith RB, Tarnow DP. Classification of molar extraction sites for immediate dental implant placement: technical note. *Int J Oral Maxillofac Implants* 2013;28:911–6.
- Tarnow DP, Chu SJ, Salama MA, et al. Flapless postextraction socket implant placement in the esthetic zone: part 1. The effect of bone grafting and/or provisional restoration on facial-palatal ridge dimensional change—a retrospective cohort study. *Int J Periodontics Restor Dent* 2014;34:323–31.
- Chappuis V, Engel O, Reyes M, et al. Ridge alterations post-extraction in the esthetic zone: a 3D analysis with CBCT. *J Dent Res* 2013;92:1955–2015.
- Ketabi M, Deporter D, Atenafu EG. A systematic review of outcomes following immediate molar implant placement based on recently published studies. *Clin Implant Dent Relat Res* 2016;18:1084–94.
- Chen YW, Finkelman M, Paspapirisdakos P, César-Neto JB, Weber HP, de Souza AB. Comparative analysis of dimensional alterations following extraction of maxillary molars using three-dimensional images superimposition: a CBCT study. *Odontology* 2021;109:514. 23.
- Ragucci GM, Elnayef B, Criado-Cámara E, Del Amo FS, Hernández- Alfaro F. Immediate implant placement in molar extraction sockets: a systematic review and meta-analysis. *Int. J. Implant Dent* 2020;6:40–51.
- Buser D, Chappuis V, Belser UC, et al. Implant placement post extraction in esthetic single tooth sites: when immediate, when early, when late. *Periodontol* 2000 2017;73:84–102.
- Lekholm U, Zarb GA. Patient selection and preparation. In: Brånemark PI, Zarb GA, Albrektsson T, eds. *Tissue-integrated prostheses: osseointegration in clinical dentistry*; 1985: 199–209. Chicago: Quintessence.
- Hämmerle CH, Chen ST, Wilson Jr TG. Consensus statements and recommended clinical procedures regarding the placement of implants in extraction sockets. *Int J Oral Maxillofac Implants* 2004;19(Suppl):26–8.
- Mustakim KR, Eo MY, Lee JY, Myoung H, Seo MH, Kim SM. Guidance and rationale for the immediate implant placement in the maxillary molar. *J Korean Assoc Oral Maxillofac Surg* 2023;49:30–42.

18. Jung RE, Zembic A, Pjetursson BE, et al. Systematic review of the survival rate and the incidence of biological, technical, and aesthetic complications of single crowns on implants reported in longitudinal studies with a mean follow-up of 5 years. *Clin Oral Implants Res* 2012;23(Suppl 6):2–21.
19. Moraschini V, Poubel LA, Ferreira VF, et al. Evaluation of survival and success rates of dental implants reported in longitudinal studies with a follow-up period of at least 10 years: a systematic review. *Int J Oral Maxillofac Surg* 2015;44:377–88.
20. Benic GI, Mokti M, Chen CJ, et al. Dimensions of buccal bone and mucosa at immediately placed implants in the anterior region: a cone-beam computed tomography study. *Clin Oral Implants Res* 2012;23:560–6.
21. Araújo MG, Lindhe J. Dimensional ridge alterations following tooth extraction. An experimental study in the dog. *J Clin Periodontol* 2005;32:212–8.
22. Wagenberg B, Froum SJ. A retrospective study of 1925 consecutively placed immediate implants from 1988 to 2004. *Int J Oral Maxillofac Implants* 2006;21:71–80.
23. Peñarrocha-Diago MA, Maestre-Ferrín L, Demarchi CL, et al. Immediate versus non-immediate placement of implants for full-arch fixed restorations: a preliminary study. *J Oral Maxillofac Surg* 2011;69:154–9.
24. Misch CE, Perel ML, Wang HL, et al. Implant success, survival, and failure: the International Congress of Oral Implantologists (ICOI) Pisa consensus conference. *Implant Dent* 2008;17:5–15.
25. Trombelli L, Farina R, Marzola A, et al. Modeling and remodeling of human extraction sockets. *J Clin Periodontol* 2008;35: 630–9.
26. Van der Weijden F, Dell'Acqua F, Slot DE. Alveolar bone dimensional changes of post-extraction sockets in humans: a systematic review. *J Clin Periodontol* 2009;36:1048–58.
27. Tufekcioglu S, Delilbasi C, Gurler G, Dilaver E, Ozer N. Is 2 mm a safe distance from the inferior alveolar canal to avoid neurosensory complications in implant surgery? *Niger J Clin Pract* 2017;20:274–7.
28. Kan JY, Rungcharassaeng K, Lozada JL, et al. Facial gingival tissue stability following immediate placement and provisionalization of maxillary anterior single implants: a 2- to 8-year follow-up. *Int J Oral Maxillofac Implants* 2011;26:179–87.
29. Evans CD, Chen ST. Esthetic outcomes of immediate implant placements. *Clin Oral Implants Res* 2008;19:73–80.
30. Pjetursson BE, Karoussis I, Bürgin W, et al. Patients' satisfaction following implant therapy. a 10 year prospective cohort study. *Clin Oral Implants Res* 2005;16:185–93.
31. Galindo-Moreno P, Catena A, Pérez-Sayáns M, Fernández-Barbero JE, O'Valle F, Padiál-Molina M. Early marginal bone loss around dental implants to define success in implant dentistry: a retrospective study. *Clin Implant Dent Relat Res* 2022;24:630–42.
32. Lombardi T, Berton F, Salgarello S, et al. Factors influencing early marginal bone loss around dental implants positioned subcrestally : a multicenter prospective clinical study. *J Clin Med* 2019;8:1168–80.
33. Karthik K, Sivakumar Sivaraj, Thangaswamy V. Evaluation of implant success: a review of past and present concepts. *J Pharm BioAllied Sci* 2013;5:5117–9.
34. Del Fabbro M, Ceresoli V, Taschieri S, et al. Immediate loading of postextraction implants in the esthetic area: systematic review of the literature. *Clin Implant Dent Relat Res* 2015;17: 52–70.
35. Chrcanovic BR, Albrektsson T, Wennerberg A. Immediately loaded non-submerged versus delayed loaded submerged dental implants: a meta-analysis. *Int J Oral Maxillofac Surg* 2015;44:493–506.
36. Witonkitvanich P, Amornsettachai P, Panyayong W, Rokaya D, Vongsirichat N, Suphangul S. Comparison of the stability of immediate dental implant placement in fresh molar extraction sockets in the maxilla and mandible: a controlled, prospective, non-randomized clinical trial. *Int J Oral Maxillofac Surg* 2025; 54:365–73.
37. Mazzocco F, Jimenez D, Barallat L, et al. Bone volume changes after immediate implant placement with or without flap elevation. *Clin Oral Implants Res* 2017;28:495–501.
38. Deporter D, Ketabi M. Guidelines for optimizing outcomes with immediate molar implant placement. *J Periodontal Implant Dent* 2017;9:37–44.
39. Becker W, Clokie C, Sennerby L, et al. Histologic findings after implantation and evaluation of different grafting materials and titanium micro screws into extraction sockets: case reports. *J Periodontol* 1998;69:414–21.
40. Botticelli D, Berglundh T, Lindhe J. Hard-tissue alterations following immediate implant placement in extraction sites. *J Clin Periodontol* 2004;31:820–8.
41. Nevins M, Camelo M, De Paoli S, et al. A study of the fate of the buccal wall of extraction sockets of teeth with prominent roots. *Int J Periodontics Restor Dent* 2006;26:19–29.
42. Quan S, Yang H, Na H, Yi J, Tong Z, Juncheng W. Restoration-oriented anatomical analysis of alveolar bone at mandibular first molars and implications for immediate implant placement surgery: a CBCT study. *J Adv Prosthodont* 2024;16:212–20.
43. Carpentieri J, Greenstein G. Guidelines for immediate Vs delayed dental implant placement in the esthetic zone. *Comp Cont Educ Dent* 2024;45:340–7.
44. Dastgardi ME, Deporter D, Xia M, Ketabi M. CBCT data relevant in treatment planning for immediate mandibular molar implant placement. *J Adv Periodontol Implant Dent* 2025;17:71–6.
45. Kato S, Kato G. Classification of molar extraction sites for immediate dental implant placement revisited: additional technical note. *J Surg* 2025;10:11479–88.
46. Aldahlawi S, Nourah DM, Azab RY, et al. Cone-Beam Computed Tomography (CBCT)-based assessment of the alveolar bone anatomy of the maxillary and mandibular molars: implication for immediate implant placement. *Cureus* 2023;15:e41608-19.
47. Kong ZL, Wang GG, Liu XY, Ye ZY, Xu DQ, Ding X. Influence of bone anatomical morphology of mandibular molars on dental implant based on CBCT. *BMC Oral Health* 2021;21:528–38.
48. Liu G, Liu R. Maxillary immediate molar implant placement into palatal root socket: a case report of two-year follow-up. *Cureus* 2024;16:e71152-6.
49. Wychowski P, Woliński J, Kacprzak M, et al. Immediate palatal molar implants: a simple, safe, minimal invasive technique. *Int J Periodontics Restor Dent* 2017;37:e297–301.
50. Bleyan S, Gaspar J, Huwais S, et al. Molar septum expansion with osseodensification for immediate implant placement, retrospective multicenter study with up-to-5-year follow-up, introducing a new molar socket classification. *J Funct Biomater* 2021;12:66–80.
51. Galindo-Moreno P, Catena A, Pérez-Sayáns M, Fernández-Barbero JE, Francisco O, Valle F, Padiál-Molina M. Early marginal bone loss around dental implants to define success in implant dentistry: a retrospective study. *Clin Implant Dent Relat Res* 2022;24:630–42.
52. Velasco-Ortega E, Wojtovicz E, España-Lopez A, et al. Survival rates and bone loss after immediate loading of implants in fresh extraction sockets (single gaps): a clinical prospective study with 4 year follow-up. *Med Oral Patol Oral Cir Bucal* 2018;23:e230–6.
53. Kim YM, Lee JB, Um HS, Chang BS, Lee JK. Long-term effect of implant-abutment connection type on marginal bone loss and survival of dental implants. *J Periodontal Implant Sci* 2022;52: 496–508.
54. Vasiljevic M, Selakovic D, Rosic G, et al. Anatomical factors of the anterior and posterior maxilla affecting immediate implant placement based on cone beam computed tomography analysis: a narrative review. *Diagnostics* 2024;14:1697–714.

55. Xiao YJ, Lv LF, Xu ZH, et al. Correlation between peri-implant bone mineral density and primary implant stability based on artificial intelligence classification. *Sci Rep* 2024;14:3009–19.
56. Wang SH, Hsu JT, Fuh LJ, Peng SL, Huang HL, Tsai MT. New classification for bone type at dental implant sites: a dental computed tomography study. *BMC Oral Health* 2023;23:324–36.
57. Cui ZM, Fang Y, Mei LZJ, et al. A fully automatic AI system for tooth and alveolar bone segmentation from cone-beam CT images. *Nat Commun* 2022;13:2096–106.